



AFRL-RX-TY-TP-2010-0099

**HIGH ELECTROCATALYTIC ACTIVITY OF  
TETHERED MULTICOPPER OXIDASE-CARBON  
NANOTUBE CONJUGATES  
(POSTPRINT)**

---

Ramaraja P. Ramasamy and Heather R. Luckarift  
Universal Technology Corporation  
1270 North Fairfield Road  
Dayton, OH 45432

Dmitri M. Ivnitski and Plamen B. Atanassov  
Department of Chemical and Nuclear Engineering  
and Center for Emerging Energy Technology  
University of New Mexico  
Albuquerque, NM 87131

Glenn R. Johnson  
Airbase Technologies Division  
Air Force Research Laboratory  
139 Barnes Drive  
Tyndall Air Force Base, FL 32403-5323

Contract No. FA4819-07-D-0001

December 2010

<b>DISTRIBUTION A:</b> Approved for public release; distribution unlimited.
---

**AIR FORCE RESEARCH LABORATORY  
MATERIALS AND MANUFACTURING DIRECTORATE**

■ Air Force Materiel Command   ■ United States Air Force   ■ Tyndall Air Force Base, FL 32403-5323

<b>REPORT DOCUMENTATION PAGE</b>				Form Approved OMB No. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p><b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b></p>					
<b>1. REPORT DATE (DD-MM-YYYY)</b> 31-DEC-2010		<b>2. REPORT TYPE</b> Journal Article POSTPRINT		<b>3. DATES COVERED (From - To)</b> 01-JAN-2008 -- 31-MAR-2010	
<b>4. TITLE AND SUBTITLE</b> High Electrocatalytic Activity of Tethered Multicopper Oxidase-Carbon Nanotube Conjugates (POSTPRINT)				<b>5a. CONTRACT NUMBER</b> FA4819-07-D-0001	
				<b>5b. GRANT NUMBER</b>	
				<b>5c. PROGRAM ELEMENT NUMBER</b> 0602102F	
<b>6. AUTHOR(S)</b> *Ramasamy, Ramaraja P.; *Luckarift, Heather R.; **Ivnitski, Dmitri M.; **Atanassov, Plamen B.; #Johnson, Glenn R.				<b>5d. PROJECT NUMBER</b> 4915	
				<b>5e. TASK NUMBER</b> L2	
				<b>5f. WORK UNIT NUMBER</b> Q140LA62	
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> *Universal Technology Corporation, 1270 N. Fairfield Road, Dayton, OH 45432 **Department of Chemical and Nuclear Engineering & Center for Emerging Energy Technology, University of New Mexico, Albuquerque, NM 87131				<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>	
<b>9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> #Air Force Research Laboratory Materials and Manufacturing Directorate Airbase Technologies Division 139 Barnes Drive, Suite 2 Tyndall Air Force Base, FL 32403-5323				<b>10. SPONSOR/MONITOR'S ACRONYM(S)</b> AFRL/RXQL	
				<b>11. SPONSOR/MONITOR'S REPORT NUMBER(S)</b> AFRL-RX-TY-TP-2010-0099	
<b>12. DISTRIBUTION/AVAILABILITY STATEMENT</b> Distribution Statement A: Approved for public release; distribution unlimited.					
<b>13. SUPPLEMENTARY NOTES</b> Ref AFRL/RXQ Public Affairs Case #10-068. Document contains color images. Published in Chem. Commun., 2010, 46, 6045-6047.					
<b>14. ABSTRACT</b> Multicopper oxidases linked to multiwall carbon nanotubes via the molecular tethering reagent, 1-pyrenebutanoic acid, succinimyd ester, displayed high bioelectrocatalytic activity for oxygen reduction.					
<b>15. SUBJECT TERMS</b> carbon nanotubes, laccase, pyrenebutanoic acid-succinimidyl ester, molecular tether, biofuel cell, multicopper oxidase					
<b>16. SECURITY CLASSIFICATION OF:</b>			<b>17. LIMITATION OF ABSTRACT</b> UU	<b>18. NUMBER OF PAGES</b> 6	<b>19a. NAME OF RESPONSIBLE PERSON</b> Glenn R. Johnson
<b>a. REPORT</b> U	<b>b. ABSTRACT</b> U	<b>c. THIS PAGE</b> U			<b>19b. TELEPHONE NUMBER (Include area code)</b>


Reset

# ChemComm

Chemical Communications

[www.rsc.org/chemcomm](http://www.rsc.org/chemcomm)

Volume 46 | Number 33 | 7 September 2010 | Pages 5977–6188



ISSN 1359-7345

RSC Publishing

**COMMUNICATION**

Glenn R. Johnson *et al.*  
High electrocatalytic activity  
of tethered multicopper  
oxidase–carbon nanotube  
conjugates

**COMMUNICATION**

Glenn R. Johnson *et al.*  
Standardized microbial fuel cell  
anodes of silica-immobilized  
*Shewanella oneidensis*



# High electrocatalytic activity of tethered multicopper oxidase–carbon nanotube conjugates

Ramaraja P. Ramasamy,<sup>ab</sup> Heather R. Luckarift,<sup>ab</sup> Dmitri M. Ivnitski,<sup>c</sup> Plamen B. Atanasov<sup>c</sup> and Glenn R. Johnson<sup>\*a</sup>

Received 13th April 2010, Accepted 4th June 2010

DOI: 10.1039/c0cc00911c


**Multicopper oxidases linked to multiwall carbon nanotubes via the molecular tethering reagent, 1-pyrenebutanoic acid, succinimidyl ester, displayed high bioelectrocatalytic activity for oxygen reduction.**

The multicopper oxidase (MCO) enzymes catalyze oxygen reduction at high electrochemical potentials, making them attractive catalysts for biological fuel cell cathodes.<sup>1</sup> The redox active center of an MCO comprises four coordinated copper atoms: type 1 (T1), type 2 (T2) and type 3 (T3) copper sites.<sup>2</sup> The T1 site lies close to the substrate-binding pocket of the enzyme. In a catalytically active system, the T1 site is the initial electron acceptor, which then subsequently passes electrons to the tri-nuclear copper center (T2 and two T3) where oxygen is ultimately reduced to water. Various experimental strategies have been proposed for establishing direct electron transfer (DET) with enzymatic electrodes in which the electrons may ‘hop’ between the enzyme redox center and the electrode. The DET mode eliminates the inherent limitations of redox mediators in biological fuel cell and biosensor applications.<sup>3</sup> The strategies to electronically connect the electrode and biocatalysts include: simple physisorption or covalent linkage, entrapment in conductive polymeric films, association with metal colloids, and encapsulation within porous matrices that incorporate a conductive nanomaterial.<sup>4</sup> Immobilization methods using carbon nanotubes (CNT) are an attractive means to create a three-dimensional, porous, conductive catalytic matrix on an electrode surface.<sup>5</sup> Establishing a methodology that will reliably link the redox enzymes to well dispersed, conductive CNT may advance bioelectrocatalysis and materials applications.

Chen *et al.* reported that CNT can be modified using 1-pyrenebutanoic acid, succinimidyl ester (PBSE). The aromatic pyrenyl moiety interacts with the aromatic-like structure of the CNT walls through irreversible  $\pi$ – $\pi$  stacking at the CNT and PBSE interface.<sup>6</sup> Subsequent incubation of the PBSE-modified CNT with protein allows the amines on the protein surface to form covalent amide bonds that link the proteins to PBSE and therefore to CNT. The PBSE-based method for sidewall

functionalization and protein immobilization on CNT has been studied with various model enzymes, but little work has been related to bioelectrocatalysis and in particular, DET of redox enzymes.<sup>7</sup>

For an initial proof of concept, two MCO were selected as model redox enzymes, laccase from *Trametes versicolor* and bilirubin oxidase (BOx) from *Myrothecium verrucaria*. The enzymes were immobilized on multi-walled CNT (MWCNT) by PBSE-modification of MWCNT directly on Toray<sup>®</sup> carbon paper (TP) electrodes (Scheme 1).<sup>†</sup> The enzyme immobilization efficiency was examined first using standard biochemical assays.<sup>†</sup> The assay showed that  $\sim 20$  U of laccase was associated with the electrode, corresponding to  $\sim 0.03$   $\mu\text{g}$  protein  $\text{mm}^{-2}$  electrode. BOx activity proved to be too low to measure spectrophotometrically for the corresponding BOx-modified PBSE/MWCNT electrode. Since the activity measured in the standard assay does not represent electrochemical activity or oxygen reduction, further electrochemical characterization of the materials was done to determine whether an electronic connection was established for the protein, MWCNT, and TP composite material.<sup>†</sup>



**Scheme 1** Illustration of MCO immobilization onto PBSE-modified carbon nanotubes.

<sup>a</sup> Microbiology and Applied Biochemistry, Air Force Research Laboratory, Tyndall Air Force Base, FL 32403, USA.  
E-mail: glenn.johnson@tyndall.af.mil; Fax: +1 850-2836090;  
Tel: +1 850-2836223

<sup>b</sup> Universal Technology Corporation, 1270 N. Fairfield Rd, Dayton, OH 45432, USA

<sup>c</sup> Department of Chemical and Nuclear Engineering & Center for Emerging Energy Technology, University of New Mexico, Albuquerque, NM 87131, USA


Cyclic voltammetry (CV) was used to evaluate the effectiveness of the PBSE-tether for functionalizing the TP/MWCNT-electrodes with enzyme. For control experiments, enzymes were associated directly to (i) TP or (ii) TP/MWCNT through noncovalent physisorption. Laccase adsorbed directly on TP showed no evidence of electrocatalytic activity for oxygen reduction (Fig. 1a). Despite a theoretically short electron tunneling distance between the enzyme redox center and the electrode, the open circuit potential (OCP) for the oxygen reduction reaction was  $0.44 \pm 0.03$  V;  $n = 4$ , significantly lower than the thermodynamic maximum ( $0.688$  V vs. Ag/AgCl) at pH 5.8. TP modified with MWCNT had higher capacitance (increased electrochemical surface area) but marginally increased OCP ( $0.49 \pm 0.02$  V;  $n = 3$ ). Although the nanomaterial dimensions may bring about close physical binding between laccase and MWCNT that could facilitate electron tunnelling, there was no evidence for interfacial DET or electrocatalytic activity.

When laccase was tethered to MWCNT *via* PBSE, the CV results depicted obvious electrocatalytic activity for oxygen reduction. The cathodic sweep showed a dramatic deflection from the control electrode processes below  $0.6$  V. The OCP, onset and half-peak potentials were  $0.60 \pm 0.01$  V,  $0.60 \pm 0.01$  V,


and  $0.47 \pm 0.02$  V;  $n = 3$ , respectively, and diffusion limitation conditions were reached at  $\sim 0.4$  V during the cathodic sweep. The voltammetric response of the PBSE-tethered laccase provided a Tafel slope of  $18$  mV per decade in the kinetic region above  $0.57$  V and a slope of  $24$  mV per decade from  $0.5$  to  $0.55$  V. The Tafel slopes approach the theoretical limit ( $15$  mV) for a four-electron-transfer reaction, and compare favorably with slopes for conventional oxygen reduction catalysts.<sup>8</sup> The CV trace for nitrogen-sparged electrolyte shows no catalytic current with the tethered laccase, confirming that the deflection seen in the cathodic sweep corresponds to oxygen reduction (Fig. 1).

The methodology was examined further using BOx as an oxygen reduction catalyst. The PBSE-tethered BOx electrode (Fig. 1b) also showed high electrocatalytic activity (OCP:  $0.62 \pm 0.005$  V;  $n = 3$ ), the onset and half-peak potentials for the electrodes were  $0.61 \pm 0.02$  V and  $0.45 \pm 0.03$  V, respectively. The OCP at neutral pH was  $\sim 0.56$  V, about  $75$  mV higher than that reported in the literature for a covalently linked BOx on MWCNT.<sup>9</sup> The CV traces of oxygen- and nitrogen-sparged electrolyte show a clear distinction between catalytic and capacitive processes, indicating a high electrocatalytic activity of BOx. Unlike laccase, BOx exhibited apparent electrocatalytic activity after simple physisorption onto the TP or TP/MWCNT electrodes without the PBSE tether. The response may result from undefined surface characteristics of BOx that bring about preferential orientation of the enzyme T1 copper site compared to laccase, or simply a higher specific catalytic activity for the commercial BOx preparation.

Galvanostatic measurements at steady state revealed that the tethered MCO exhibit exceptionally stable performance with potential losses of less than  $100$  mV at  $50 \mu\text{A cm}^{-2}$ , relative to the OCP. By comparison, laccase physisorbed onto TP or TP/MWCNT-modified could not sustain high faradic currents (Fig. 2). The BOx on TP and TP/MWCNT



**Fig. 1** Cyclic voltammograms (CV) of model MCO bioelectrodes (laccase (a), BOx (b)). Key: (1) MCO physisorbed on bare TP, (2) MCO physisorbed on TP/MWCNT electrode; (3) MCO immobilized on PBSE-modified TP/MWCNT electrode; (4) electrode 3 in nitrogen-flushed electrolyte. CV scans in phosphate buffer electrolyte (pH 5.8), scan rate  $10 \text{ mV s}^{-1}$ , oxygen saturated electrolyte except as noted.



**Fig. 2** Galvanostatic polarization curves for laccase (solid lines, closed symbols) and BOx (dotted lines, open symbols) electrodes. Key: (triangles) MCO physisorbed on bare TP; (circles) MCO physisorbed on TP/MWCNT electrode; (squares) MCO immobilized on PBSE-modified TP/MWCNT electrode. Note: potentials  $< 0$  are not shown; the plotted curves extrapolate beyond the x-axis and are based on all measured data.

electrodes, however, showed comparably better performance than laccase controls, in agreement with the CV results.

The PBSE-modified TP/MWCNT electrodes supported faster and more complete bioelectrochemical oxygen reduction than the physisorbed control materials and facilitated DET for immobilized MCO at current densities much greater than previous reports.<sup>9</sup> The galvanostatic measurements for the MCO confirm the fabrication of a stable, conductive, bioelectrocatalytic interface between the enzyme and electrode that may be attributed to a variety of factors. The PBSE-modification of MWCNT *via*  $\pi$ - $\pi$  stacking preserves the electronic properties of the MWCNT to allow efficient electron transport through the matrix. The covalent link between a MCO and the PBSE-tether will position the enzyme close to the MWCNT and reduce the electron tunneling distance between the enzyme and the electrode. In addition, the distribution of amines on the protein surface will guide orientation with the MWCNT. The limited electrochemical activity measured for control electrodes (no PBSE) is attributed to a small fraction of MCO molecules that align favorably during physisorption on TP or MWCNT. The response is evident from the higher OCP measured with MWCNT on TP (0.49 V) when compared to that of protein-free MWCNT (0.09 V). The high OCP, however, did not translate to electrocatalytic activity, possibly due to unfavorable orientation of the majority of catalyst molecules on the MWCNT surface or poor catalyst loading. A detailed understanding of the biophysical interactions at the interface that lead to efficient bioelectrocatalysis will require additional characterization and modeling.

The bio-conjugates formed using PBSE effectively link MCO with MWCNT to facilitate DET and bioelectrocatalytic oxygen reduction. The catalytic efficiency was significantly greater than previous reports for MCO electrodes. The process provides a porous, potentially scalable, architecture that can advance bioelectrocatalytic applications. Future research will provide a deeper understanding of the attachment mechanism that directs enzyme orientation and provides guidance to optimize the interaction further. The applicability to alternative catalysts was demonstrated with two MCOs but could be extended to a wider range of biomolecules and applications.

AFRL work was supported by the AFOSR and AFRL-Materials and Manufacturing Directorate. The UNM research was supported by DOD/AFOSR MURI (code FA9550-06-1-0264).

## Notes and references

† Preparation of tethered MCO-MWCNT conjugates: 10 mm<sup>2</sup> discs of Toray<sup>®</sup> carbon paper (TGP 090; Toray Industries, Tokyo, Japan) used as electrode base. For CNT modification of TP, 30  $\mu$ L of MWCNT (1 mg mL<sup>-1</sup> in *N,N*-dimethyl formamide) (Sigma, St. Louis, MO) was drop-cast onto TP and dried (50 °C, 30 min). For sidewall functionalization of CNT with PBSE, the TP/MWCNT electrodes were soaked in a PBSE solution [1-pyrenebutanoic acid succinimidyl

ester; Anaspec Inc, Fremont, CA; (10 mM in *N,N*-DMF)] for 1 h at room temperature, washed thoroughly with *N,N*-DMF to remove excess PBSE and then with phosphate buffer (10 mM, pH 7). Laccase (Sigma) was dialyzed against CuSO<sub>4</sub> (10  $\mu$ M) in phosphate buffer (20 mM, pH 5.8) and then stored at -20 °C until use. Protein concentrations were determined using BCA assay (BCA Protein Assay Kit, Thermo Scientific Inc. Rockford, IL). BOx (Sigma) was used as received. For protein immobilization, PBSE-modified MWCNT/TP electrodes were incubated with 0.2 mL of laccase (0.5 mg mL<sup>-1</sup>) or BOx (0.2 mg mL<sup>-1</sup>) for 1 h in phosphate buffer (10 mM, pH 7.0). Excess protein was removed by washing with phosphate buffer and the electrode tested immediately in an electrochemical cell. Enzyme assays: the oxidase activity of laccase was determined in phosphate buffer (100 mM, pH 6.5) using syringaldazine (21.6  $\mu$ M in methanol) (Sigma) by measuring the change in absorbance at 530 nm over time. One unit of activity was defined as 1  $\mu$ mol of syringaldazine oxidized by laccase in 1 minute at 37 °C. The catalytic activity of BOx was evaluated using bilirubin as substrate (0.002% in 0.2 M Tris HCl buffer, pH 8.4) by measuring the change in absorbance at 440 nm over time. One unit of activity was defined as 1  $\mu$ mol of bilirubin oxidized by BOx in 1 minute at 37 °C. Electrochemical testing: the electrodes were held on a capped glassy carbon shaft electrode and tested in a 50 mL voltammetric cell (CH Instruments Inc, Austin, TX) with a glassy carbon counter electrode and a Ag/AgCl reference electrode (CH Instruments Inc.). Phosphate buffer (100 mM, pH 5.8, or pH 7.0 when noted) was used as the electrolyte throughout. CV scans were obtained by scanning from -0.2 V to 0.8 V at a sweep rate of 10 mV s<sup>-1</sup>. Galvanostatic polarization measurements were obtained from 0 to 50  $\mu$ A cm<sup>-2</sup> at 10  $\mu$ A cm<sup>-2</sup> intervals and the voltage data points were recorded after 15 min at each step. Tafel slopes were obtained from the plots of working electrode potential versus kinetic current in the absence of convective transport.

- 1 S. C. Barton, H.-H. Kim, G. Binyamin, Y. Zhang and A. Heller, *J. Am. Chem. Soc.*, 2001, **123**, 5802–5803; S. C. Barton, J. Gallaway and P. Atanassov, *Chem. Rev.*, 2004, **104**, 4867–4886; W. Zheng, H. Y. Zhao, H. M. Zhou, X. X. Xu, M. H. Ding and Y. F. Zheng, *J. Solid State Electrochem.*, 2010, **14**, 249–254.
- 2 E. I. Solomon, U. M. Sundaram and T. E. Machonkin, *Chem. Rev.*, 1996, **96**, 2563–2605; K. Piontek, M. Antorini and T. Choinowski, *J. Biol. Chem.*, 2002, **277**, 37663–37669.
- 3 S. Shleev, J. Tkac, A. Christenson, T. Ruzgas, A. I. Yaropopov, J. W. Whittaker and L. Gorton, *Biosens. Bioelectron.*, 2005, **20**, 2517–2554.
- 4 L. M. Ellerby, C. R. Nishida, F. Nishida, S. A. Yamanaka, B. Dunn, J. S. Valentine and J. I. Zink, *Science*, 1992, **255**, 1113–1115; D. Ivnitiski, K. Artyushkova, R. A. Rincon, P. Atanassov, H. R. Luckarift and G. R. Johnson, *Small*, 2008, **4**, 357–364; M. Mazur, A. Krywko-Cendrowska, P. Krynski and K. Rogalski, *Synth. Met.*, 2009, **159**, 1731–1738; H. R. Luckarift, D. Ivnitiski, R. Rincon, P. Atanassov and G. R. Johnson, *Electroanalysis*, 2010, **22**, 784–792.
- 5 J. M. Guisan, *Immobilization of enzymes and cells*, Humana Press, Totowa NJ, USA, 2006; P. F. Harris, *Carbon Nanotube Science*, Cambridge Publishers, Cambridge, UK, 2009.
- 6 R. J. Chen, Y. Zhang, D. Wang and H. Dai, *J. Am. Chem. Soc.*, 2001, **123**, 3838–3839.
- 7 G. T. Hermanson, *Bioconjugate Techniques*, Elsevier Academic Press, New York NY, USA, 2008; K. Besteman, J.-O. Lee, F. G. M. Wiertz, H. A. Heering and C. Dekker, *Nano Lett.*, 2003, **3**, 727–730.
- 8 E. Gileadi, *Electrode Kinetics*, VCH Publishers, New York, NY, USA, 1993.
- 9 W. Zheng, H. M. Zhou, Y. F. Zheng and N. Wang, *Chem. Phys. Lett.*, 2008, **457**, 381–385; M. Jonsson-Niedziolka, K. Szot, J. Rogalski and M. Apallo, *Electrochem. Commun.*, 2009, **11**, 1042–1044; K. Schubert, G. Goebel and F. Lisdat, *Electrochim. Acta*, 2009, **54**, 3033–3038.